

THE NATURE OF DARK MATTER

Ben Moore

Department of Astronomy, University of California, Berkeley, CA 94720, USA

ABSTRACT

Collisionless particles, such as cold dark matter, interact only by gravity and do not have any associated length scale, therefore the dark halos of galaxies should have negligible core radii. This expectation has been supported by numerical experiments of collisionless particles within scale free and cold dark matter cosmologies. Most dwarf spiral galaxies are almost completely dark matter dominated, allowing a unique insight into their mass – density profiles which can be approximated by isothermal spheres with core radii of order 3 – 7 kpc. We can therefore make a direct comparison between these galaxies, and halos which form within the numerical simulations. This yields a severe discrepancy in that the simulations predict the density to fall as $\rho(r) \propto r^{-1}$ on the scales where the data show that $\rho(r) = \text{const}$: *e.g.* the models overestimate the mass within the central few kpc of the halos by a factor of four. The formation of the luminous component of galaxies exacerbates this disparity between theory and observations, since the contraction of the baryons can significantly increase the central dark matter density.

One of the primary motivations for invoking dark matter halos arose when radio observations of neutral hydrogen (HI) in spiral galaxies at large galactocentric radii, r_g , showed evidence for more mass than was observed within the visible components¹. In general, the gas rotates at an approximately constant velocity out to the last measured data point^{2,3}, implying that the density of material falls approximately as r_g^{-2} , *i.e.* the mass increases linearly with distance. For r_g less than the characteristic optical radius, the dynamics are dominated by the luminous material, whereas on larger scales a dark matter component is required to maintain the motion of the observed baryons. The importance of the dark halo at small radii depends upon the mass to light ratio (M/L) of the baryonic material in the disk; typically, M/L ratios of order unity are adopted. This would imply that in a large spiral galaxy like our own, the gravity of the luminous material dominates over that of the dark matter to distances of order ~ 10 kpc.

Determining the structure of dark halos from the observations is difficult, and only in a handful of galaxies is this value tightly constrained due to the uncertainties in the mass of the disk material and the unknown dark matter distribution². Maximum disk models attempt to fit the inner part of the rotation curve using only the observed bulge and disk components and adopting high M/L ratios, thus requiring extended dark halos with core radii as large as 30 kpc²⁻⁵. If very low mass to light ratios are adopted, then the core radii can be significantly reduced, but not completely. Dwarf spiral galaxies provide excellent probes of the internal structure of dark halos, since these galaxies are completely dominated by dark matter on scales larger than a few kpc⁶⁻⁸. Furthermore, most of the baryonic mass is in the form of HI within the disk, which can be directly weighed using the observed 21 cm emission.

At present less than a dozen rotation curves have been measured for dwarf galaxies, however a trend is clearly apparent in that the rotational velocity of the gas is typically rising over most of the data, many optical scale lengths into the dark halo, but within the core of the mass distribution (see Figure 1). For example, the rotation curve of the dwarf galaxy DDO 154 shown in Figure 1, is dominated by dark matter beyond about 1.5 kpc⁸. In this system the amount of HI, $2.7 \times 10^8 M_\odot$, is 5 times the stellar component and extends over 15 optical scale lengths into the dark halo which has total mass $\gtrsim 5 \times 10^9 M_\odot$. Although dwarf spirals such as the DDO galaxies have luminosities only a few percent of an L_* galaxy, these galaxies are the most numerous, and a large fraction of the mass in the Universe is associated with these systems. For comparison, the rotation curve of NGC 925 is plotted in Figure 1⁹. This is a normal spiral galaxy, but even this system has a dark halo with large core radius, $r_c \gtrsim 7$ kpc.

Any successful cosmological model must be able to reproduce both the observed large and small scale structures, from galaxy clusters to galaxy halos. A Universe dominated by collisionless particles, such as axions or gravitinos which interact only through gravity¹⁰, has been extensively investigated^{11–14}. One of the major successes of the so called cold dark matter (CDM) models, is the formation of dark halos whose density profiles fall approximately as r_g^{-2} ; implying a circular velocity, $v_c(r_g) = (Gm/r_g)^{1/2}$, which is constant with distance^{15,16}. The extent of the dark halos in these models depends on Ω : If $\Omega \approx 0.1$, then the halo of our Galaxy would extend to touch that of Andromeda's at ≈ 300 kpc.

The inner structure of halos which form within CDM and scale free Universes dominated by collisionless particles, has recently been re-examined at a resolution 50 times higher than the original simulations^{17,18}. This work confirms the original numerical investigations^{15,16}, and concludes that halos form with nearly flat rotation curves, with small gradients which depend upon the slope of the initial power spectrum. *Moreover, the density continues to increase to distances less than the resolution limit of ~ 1 kpc, and the core radius, if present, must be smaller than the resolution scale.* These simulations lend strong support to the theoretical expectation that the clustering of collisionless dark matter particles is scale free, and dark halos do not have core radii.

The density profiles of the simulated halos can be fit relatively well using a Hernquist profile¹⁷, where on scales between $10 \sim 50$ kpc the density falls roughly as r_g^{-2} , and on small scales as r_g^{-1} . The resulting rotation curve, $v_c(r_g) = (Gm_s r_g)^{1/2} / (r_g + r_s)$, is parameterised by the effective mass, m_s , which can be taken as the mass within the virial radius, and r_s is a scale representing the peak height of the initial density fluctuation¹⁷. Dubinski & Carlberg¹⁷ simulated the formation of a large halo with peak circular velocity $v_{c,max} \approx 280$ km s⁻¹, and obtained values of $m_s = 2.0 \times 10^{12} M_\odot$ and $r_s = 26$ kpc. Warren *et al.*¹⁸ present the rotation curve of a halo from one of their simulations, which has a peak circular velocity $v_{c,max} \approx 140$ km s⁻¹. For this halo the Hernquist profile with parameters $m_s = 2.5 \times 10^{11} M_\odot$ and $r_s = 13$ kpc fits very well on scales up to about 20 kpc (although it appears to fall about 20% too low on larger scales). These parameters agree well with those obtained by Dubinski & Carlberg¹⁷, scaled to the virial mass of the halo, *i.e.* $m_s \propto v_{c,max}^3$, and assuming that $r_s \propto v_{c,max}$. (The rotation curves presented by Warren *et al.*¹⁸ appear to demonstrate that this scaling holds down to the smallest halos in their simulations which have amplitude ≈ 60 km s⁻¹.)

Figure 1 shows the predicted rotation curves for several spiral galaxies obtained using Hernquist profiles with r_s scaled as above, and m_s adjusted so that the curves attain

the observed $v_{c,max}$. The disparity between the models and the data follows naturally from the nature of collisionless particles which have no physically associated length scale, therefore dark matter halos should have zero effective core radii. This applies to any model in which the dark matter behaves predominantly like CDM, *e.g.* mixed dark matter models¹⁹, CDM models with a cosmological constant²⁰, or baryonic isocurvature fluctuation models which form compact dark objects very early²¹.

CDM particles are by definition moving slowly at the present epoch, therefore unlike hot dark matter particles (HDM), such as light neutrinos, primordial phase space constraints do not impose a cosmologically significant scale. Although HDM has been excluded from the dark matter candidate list for other reasons^{22–24}, phase space constraints do provide an attractive mechanism to limit the central density of dark matter particles. An additional phase space constraint might arise through the merging of halos which tends to populate the available coarse grained phase space, reducing a pre-existing core until phase space is filled²⁵. Subsequent merging could possibly reduce the central density of dark matter particles, but the cosmological simulations discussed above^{17,18} demonstrate that if this effect occurs, then it is on a scale smaller than the resolution limit. Numerical effects, such as two body relaxation, also appear to be unimportant, since identical results were obtained after increasing the numbers of particles by an order of magnitude, and using different N-body codes.

The numerical simulations discussed above only follow a dissipationless component, however gas cools efficiently within dark halos²⁶ and can modify the halo mass distribution. For example, the adiabatic cooling of gas which contracts to form the bulge and disk, *increases* the central density of dark matter and would tend to erase any pre-existing core radii^{27,28}. For a baryonic fraction of 10%, the final core radius of a dark halo may have contracted by up to a factor of three after the baryons have cooled²⁷, thus making the case against exotic dark matter even stronger. The rotation curves of bright compact spiral galaxies appear to support this notion, in that these systems have falling rotation curves due to the large amount of dissipation which has occurred, as compared with dwarf galaxies or normal spirals which have rising or flat rotation curves²⁹. Even 100% efficient energy transfer (in the opposite sense as discussed above - assuming such a mechanism exists) between the baryons which could cool within a Hubble time and the dark matter halos, would be insufficient to imprint a scale the size of the observed core radii.

Baryonic dark matter has recently achieved renewed popularity, and its nature is relatively well constrained; low mass stars remain one of the strongest candidates³⁰.

Baryons can interact via electromagnetic forces as well as gravity and therefore have the potential to introduce a characteristic scale, perhaps resulting from supernovae driven winds or cooling flows. The smooth transition between the disk material and halo material in large spiral galaxies, and the constant ratio of HI surface density to the dark matter surface density within dwarf galaxies, represent the well known ‘disk – halo’ conspiracies^{1,8}, which are easily explained if the dark matter is itself baryonic.

We have focussed on the halos of dwarf spiral galaxies whose tractable density profiles clearly hold important clues as to the nature of dark matter and the process of galaxy formation. The observations are inconsistent with the dominant dark matter component being a collisionless, weakly interacting particle such as invoked by the various CDM scenarios. A candidate particle with primordial phase space constraints does not exist, therefore unless the numerical simulations are giving misleading results on scales larger than 5 times the resolution limit, we are drawn to the conclusion that the dark matter consists of baryons.

Acknowledgements I would like to thank Marc Davis for many useful and interesting discussions.

REFERENCES

1. Faber S.M. & Gallagher J.S. 1979, *Ann.Rev.Ast.Astr.*, **17**, 135-87.
2. Lake G. & Feinswog L. 1989, *A.J.*, **98**, 166-79.
3. Begeman K.G. 1989, *A.A.*, **223**, 47-60.
4. Athanassoula E., Bosma A. & Papaioannou S. 1987, *A.A.*, **179**, 23-40.
5. Kent S.M. 1987, *A.J.*, **93**, 816-32.
6. Jobin M. & Carignan C. 1990, *A.J.*, **100**, 648-62.
7. Lake G., Schommer R.A. & van Gorkom J.H. 1990, *A.J.*, **99**, 547-60.
8. Carignan C. & Beaulieu S. 1989, *Ap.J.*, **347**, 760-70.
9. Wevers B.M.H.R. 1984, *Ph.D. thesis*, University of Groningen.
10. Sarkar S. 1991, *NATO A.S.I.*, **348**, 91-102.
11. Davis M., Efstathiou G., Frenk C.S. & White S.D.M. 1985, *Ap.J.*, **292**, 371-94.
12. White S.D.M., Frenk C.S., Davis M. & Efstathiou G. 1987, *Ap.J.*, **313**, 505-16.
13. Weinberg D.H. & Cole S. 1992, *M.N.R.A.S.*, **259**, 652-94.
14. Moore B., Frenk C.S. & White S.D.M. 1993, *M.N.R.A.S.*, **261**, 827-46.
15. Frenk C.S., White S.D.M., Efstathiou G.P. & Davis M. 1985, *Nature*, **317**, 595-7.
16. Quinn P.J., Salmon J.K & Zurek W.H., 1986, *Nature*, **322**, 329-35.
17. Dubinski J. & Carlberg R. 1991, *Ap.J.*, **378**, 496-503.
18. Warren S.W., Quinn P.J., Salmon J.K. & Zurek H.W. 1992, *Ap.J.*, **399**, 405-25.
19. Davis M., Summers F.J. & Schlegel D. 1992, *Nature*, **359**, 393-96.
20. Efstathiou G.P., Bond J.R. & White S.D.M. 1992, *M.N.R.A.S.*, **258**, 1-6P.
21. Peebles P.J.E. 1987, *Nature*, **327**, 210-11.
22. White S.D.M., Davis M. & Frenk C.S. 1984, *M.N.R.A.S.*, **209**, 27-31P.
23. Spergel D.N. 1990, *Nuclear Phys. B.*, **13**, 66-74.
24. Gerhard O.E. 1992, *Ap.J.Lett.*, **389**, L9-11.
25. Pearce F.R., Thomas P.A. & Couchman H.M.P. 1993, *M.N.R.A.S.*, **264**, 497-508.
26. Katz N., Hernquist L. & Weinberg D.H. 1992, *Ap.J.Lett.*, **399**, L109-12.
27. Blumenthal G., Faber S., Flores G. & Primack J. 1986, *Ap.J.*, **301**, 27-34.
28. Carlberg R.G., Lake G. & Norman C.A. 1986, *Ap.J.Lett.*, **300**, L1-4.
29. Casertano S. & van Gorkom J.H. 1991, *A.J.*, **101**, 1231-41.
30. Gerhard O.E. & Silk J. 1994, *Nature* in press.

FIGURE 1

The observed rotation curves of four spiral galaxies are plotted as a function of galactocentric distance. The dotted lines show a fit to the observed rotation curve assuming zero contribution from the baryons, and a dark halo with an approximately isothermal density profile $\rho(r) \propto 1/(r_c^2 + r^2)$, where r_c is the core radius. These ‘maximum halo’ models fit the dwarf galaxies very well and demonstrate that even if the gravity of the baryons is ignored, the dark halos must have large core radii. The solid curves, plotted alongside the data for DDO 154 and NGC 3109, show the contribution of the dark halo to the rotation curve after including the contribution of the stellar and HI mass. The core radii of these dark halos are fairly tightly constrained and show an interesting trend of increasing size with $v_{c,max}$. (References for the data: DDO 154⁸, DDO 170⁷, NGC 3109⁶, NGC 925⁹.) The dashed curves show the rotation curves obtained adopting a Hernquist density profile, and are drawn fainter on scales below the resolution limit of the simulations. The effective radius is scaled using $r_s \propto v_{c,max}(\text{observed})$ and the effective mass scaled so that the curve passes through this same point. (Fixing m_s this way yields curves with amplitudes about 20% lower than scaling $m_s \propto v_{c,max}^3$, where $v_{c,max}$ is obtained from the isothermal halo fit to the data.) The model predictions overestimate the observed mass within $r_c/2$ by a factor of 4 in each galaxy. The rotation curve of NGC 3109 implies that for $r < 4$ kpc, $\rho(r) \propto r^{0.0}$ to 5% accuracy, whereas the Hernquist profile predicts $\rho(r) \propto r^{-1}$. Even if r_s and m_s are left as free parameters, this profile cannot provide an acceptable fit to the rotation curves of dwarf spirals.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/astro-ph/9402009v1>